

Amendments to the Specification

Please replace the paragraph beginning at page 7, lines 1-11, with the following rewritten paragraph:

include imipramine, amitriptyline, nortriptyline, and desipramine. Monoamine oxidase inhibitors (MAOIs) are also used as antidepressants. MAOIs approved for the treatment of depression include phenelzine (~~Nardil~~ NARDIL®), tranylcypromine (PARNATE® ~~Parnate~~), and isocarboxazid (MARPLAN® ~~Marplan~~). Medications that primarily affect the neurotransmitter serotonin, termed selective serotonin reuptake inhibitors (SSRIs), are also used as antidepressants. These include escitalopram HBr (LEXAPRO® ~~Lexapro~~), fluoxetine (PROZAC® ~~Prozac~~), sertraline (ZOLOFT® ~~Zoloft~~), fluvoxamine (LUVOX® ~~Luvor~~), paroxetine (PAXIL® ~~Paxil~~), and citalopram (CELEXA® ~~Celexa~~). Additional medication of use affect both norepinephrine and serotonin, for example venlafaxine (~~Effexor~~ EFFEXOR®) and nefazadone (SERZONE® ~~Serzone~~), or agents such as phenelzine (~~Nardil~~-NARDIL®), tranylcypromine (PARNATE® ~~Parnate~~), mirtazepine (REMERON® ~~Remeron~~), nefazodone (SERZONE® ~~Serzone~~), triazolopyridine (TRAZODONE® ~~trazodone~~), and bupropion (WELLBUTRIN® ~~Wellbutrin~~).

Please replace the paragraphs on page 17, lines 5-8 with the following rewritten paragraph:

Example 1

Patient 1

A 59-year-old Caucasian female with a history of depression for 3 years was treated. Medications included PAXIL® Paxil (20 mg), as prescribed by her psychiatrist. She appeared somewhat depressed at the time of the first office visit. Her glabellar frown lines and forehead lines were treated with 40 i.u. of BOTOX™ (Fig. 4).

Please replace the paragraph at page 17, lines 22-27, with the following rewritten paragraph:

Example 2

Patient 2

A 33-year-old Caucasian female with a history of intermittent anxiety/depression for ten years was treated. Her medications included an SSRI (ZOLOFT® Zoloft, 200 mg per day) for the last four years. Prior to treatment she reported feeling depressed and anxious.

Please replace the paragraph at page 18, lines 1-10 with the following rewritten paragraph:

Example 3

Patient 3

A 50-year-old, recently divorced, Caucasian female was treated. She had a history of depression for two years. Her medications included PAXIL® Paxil-20 mg orally per day and Wellbutrin 225 mg orally per day. She reported feeling depressed.

Her frown/glabellar lines were injected with 30 Unit equivalents of BOTOX™ (Fig. 6). Four months later she reported a much-improved mood, and stated that she did not feel depressed. Her frown/glabellar lines were re-treated with 30 Unit equivalents of BOTOX™.